

The Relationship Between Overweight and Psychological Problems Among University Students in Saudi Arabia

Abdul Wahab Pathath^{1*}, Nurjahan Begum²

ABSTRACT

Background: During the last few decades, the Kingdom of Saudi Arabia (KSA) experienced rapid socio-cultural changes caused by the accelerating economy in the Arabian Gulf region. That was associated with major changes in the food choices and eating habits, which, progressively, became more and more "Westernized". Such "a nutritional transition" has been claimed for the rising rates of overweight and obesity, which were recently observed among Saudi population. **Objectives:** To identify the effect of overweight on psychological problems among university students in Saudi Arabia. **Method:** This cross-sectional study was conducted from June to November 2022 where 150 medical students were selected by convenient sample technique. Psychological problems like depressive and anxiety symptoms were measured by using DASS (Lovibond, S. & Lovibond, P. 1995; Short version) self-report questionnaire. **Results:** The findings of the study indicated significant positive relationship between overweight and psychological problems ($r = .61, p < .01$). Body mass index was found significantly and positively correlated with anxiety aspect of psychological wellbeing. **Conclusion:** It is necessary to implement adequate health promotion programs considering the high prevalence and increasing incidence of overweight in Saudi Arabia, which affect all age groups.

Keywords: *Psychological Problem, Overweight, Saudi Arabia*

Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have a negative effect on health. People are generally considered obese when their body mass index (BMI), a measurement obtained by dividing a person's weight by the square of the person's height, is over 30 kg/m^2 , with the range $25\text{--}30 \text{ kg/m}^2$ defined as overweight. Obesity increases the risk of many physical and mental conditions.

In Saudi Arabia, obesity is one of the most common chronic health problems (World Federation for Mental Health, 2010). There is presently dearth of psychological research on overweight and mental health conducted on University student population, especially in Saudi Arabia. Till date, many studies have been conducted independently on obesity and

¹Department of Clinical Neurosciences, College of Medicine, King Faisal University, Al Ahsa, 31982, Saudi Arabia

²Department of Clinical Neurosciences, College of Medicine, King Faisal University, Al Ahsa, 31982, Saudi Arabia

*Corresponding Author

Received: January 17, 2024; Revision Received: April 07, 2024; Accepted: April 12, 2024

The Relationship Between Overweight and Psychological Problems Among University Students in Saudi Arabia

mental health problems in Saudi Arabia. Overweight is associated with an increased risk of mental illness; however, evidence linking body mass index (BMI) to mental health disorder is lacking.

Obesity is defined as a condition of abnormal or excessive fat accumulation in adipose tissue, to the extent that mental health may be impaired. The most common measure used to determine relative body weight is body mass index (BMI) which is calculated as body weight in kilograms divided by the square of body height in metres (kg/m²). Categorization of individuals into groups of underweight, normal weight, overweight, and obese is commonly done using the cut-off values provided by the World Health Organization (2010). For adults 18 years or older, underweight is defined as BMI below 18.5, normal weight between 18.5 and 24.9, overweight between 25 and 29.9, and obesity as BMI of 30 or higher. The cutoff value for obesity (BMI \geq 30kg/m²) was based primarily on the reported associations between BMI and mortality. For individuals younger than 18 years, age- and sex-specific cut-off values for overweight and obesity are used.

WHO reports prevalence of excessive weight and obesity in males and females aged between 18 and 21 in the Kingdom of Saudi Arabia (KSA) is 23% and 30% respectively (Al-Othaimenet et al., 2007). According to a study, incidence of excess of weight and obesity reached 64.5% in the Eastern region, which is considered the highest in KSA (Norah M. 2010). The foremost curable cause of overweight is the dysregulation of energy consumption and expenditure, related to inappropriate dietary habits and lack of exercise (Al-Rethaiaan et al., 2010). Recent studies revealed westernization of Saudi livelihood being the main culprit in this regard. The Saudi population prefers sedentary over dynamic life style and desires burgers and pizza rather vegetables and fruits. These changes are considered to be the main reasons for the increase in the prevalence of both overweight and obesity among Saudi children, adolescents and adults in the last few decades (Mahfouz A. et al., 2007).

Many college students experience changes in their life patterns and a weakening of dietary habits during their college years. Numerous studies reported that college students have inappropriate eating habits such as skipping meals, taking high-energy intake with high fat and sodium but low calcium and iron. (Fernandes J, et al., 2013; Al-Othaimenet al., 2007; Iannotti R & Wang, J, 2013; Yaguchi-Tanaka Y, et al., 2013). According to a WHO report, there is a correlation between low Physical Activity (PA) on the one hand, and improper diet, excessive weight and obesity on the other (WHO, 2010). Al-Hazzaa et al. (2011) reported in their study on Saudi youth that 91.2% of females and 84% of males included in their study population spent more than 2hrs daily on the computer or watching TV. Almost 75% of females and 50% of the males did not do recommended daily physical activity. (Al-Hazzaa H. at al., 2011) Women in KSA are traditionally more inactive than men. It was reported that female public schools in KSA do not provide physical education to their students. (Al-Hazzaa H. at al., 2011; Khalaf A at al., 2013) The lifestyles of young adult females including their dietary patterns established during college years have a significant influence on the health of their prospective family.

LITERATURE REVIEW

The effects of overweight on mental health have been examined by different researchers and they found different results (Frisco, Houle, & Lippert, 2013; Frisco, Houle, & Martin, 2009; Needham, B. et al., 2010, Kivimäki, M. at el., 2009; Rhew et al., 2008; Richardson, Garrison,

The Relationship Between Overweight and Psychological Problems Among University Students in Saudi Arabia

Drangsholt, Mancl, & LeResche, 2006; Sjöberg, Nilsson, & Leppert, 2005). A meta-analysis pooling the results of 17 cross-sectional studies concluded that the association between overweight and depression was highly statistically significant and possibly varied by gender (de Wit, L. et al., 2010). Sjöberg, Nilsson, and Leppert (2005) investigated the relationship between overweight and depression among 4,703 adolescents in the ages of 15 to 17. Their results showed that there was a significant relationship between overweight and depression, where obese participants reported more symptoms of depression than their peers. Frisco and his associates (2013) used data from US-Based National Longitudinal study of Adolescent Health to examine what effects changes in weight during adolescence could have on depression. Their results showed that participants who became obese during adolescence had double the odds of suffering from depression during early adulthood. The results from a study conducted by Richardson and his associates were similar. They examined the association between overweight and depression during puberty among 3,101 adolescents who were 11-17 years old. Their results showed that participants who had high levels of depression were twice as likely to be obese. There is evidence that overweight and obesity are not associated with depression (Erermis et al., 2004; Mustillo et al., 2003; Swallen, Reither, Haas, & Meier, 2005; Wardle, Williamson, Johnson, & Edwards, 2006). Swallen and his associates (2005) did not find a significant relationship between overweight and obesity among 4,734 adolescents in grades 7 through 12. Neither did Wardle and his associates (2006), who twice examined the relationship between adolescent overweight and depression. Participants in the first study were 4,320 7th graders and the second study had 1,824 participants, also in the 7th grade. Results of these studies showed that there was no significant relationship between overweight and depression among participants in both samples. Needham and Crosnoe (2005) examined the association between overweight and depression among 18,924 adolescents between the ages 11 and 21 and found association between overweight and depression, but only among females. Rhew and associates (2008) also found a significant association between overweight and depression among 446 6th graders, their results showed that depressed adolescent females were more likely to be overweight and obese than non-depressed adolescent females. There is no apparent reason for the different results between researches on the association between overweight and obesity among adolescents and their symptoms of depression. The effects of overweight and obesity on symptoms of anxiety among adolescents have also been examined (Anderson, Cohen, Naumova, Jacques, & Must, 2007; Erermis et al., 2004; Eschenbeck, Kohlmann, Dudey, & Schurholz, 2009; Lanza, Echols, & Graham, 2012). Erermis and his colleagues (2004) examined the frequency of psychiatric disorders, including anxiety disorder, among obese adolescents. Participants (N=90, 12-16 years of age) were divided evenly into three groups; clinically obese, non-clinically obese and those of normal weight. The results of the study showed that participants who were obese had higher levels of anxiety compared with participants who were of normal weight. The results of a study conducted by Anderson and his associates (2007) showed that obesity could be linked to the development of anxiety disorder, but only among adolescent females. Eschenbeck and his associates (2009) also found an association between overweight or obesity and anxiety among adolescent females. It has also been shown that overweight and obese adolescent females report more symptoms of anxiety than their peers (Lanza et al., 2012). Another recent systematic review and meta-analysis found a weak but positive association between overweight and anxiety disorders. (Garipey G, 2010)

Because of the different results between researches regarding the relationship between overweight and mental health it is important to explore this further. More longitudinal

The Relationship Between Overweight and Psychological Problems Among University Students in Saudi Arabia

studies need to conduct to see the long-term effects of overweight on symptoms of depression, especially in university students. Therefore, the objective of this study is to examine the relationship between overweight and mental health among university students in Al Ahsa, Saudi Arabia.

In view of the above the following hypotheses will be tested in the present study.

- Overweight will be significantly related to the student's mental health.
- Overweight and mental health of male and female students will be significantly different.
- Demographic characteristics of participants will significantly explain variance in overweight and mental health of the participants

METHODS

Sample

A sample of 150 university students (75 male and 75 female) will be randomly selected from different colleges of King Faisal University located in Al Ahsa, Saudi Arabia. The age of these participants will be ranging from 17 to 28 years.

Measures

The following measurement instruments will be used in the present study.

- 1. Demographic questionnaire:** The information about demographic profile of the participants will be collected with the help of questions related to their age, height, weight, sex, marital status, course of study, and academic year. In addition, information about their family which include area of residence, family structure, education level of parents, family occupation, income, housing status etc.
- 2. DASS:** The DASS (Lovibond, S. & Lovibond, P. 1995; Short version) is a 21 item self report questionnaire designed to measure the severity of a range of symptoms common to both Depression and Anxiety. Each item is scored from 0 (did not apply to me at all over the last week) to 3 (applied to me very much or most of the time over the past week). The essential function of the DASS is to assess the severity of the core symptoms of Depression, Anxiety and Stress. Accordingly, the DASS allows not only a way to measure the severity of a patient's symptoms but a means by which a patient's response to treatment can also be measured. The scale to which each item belongs is indicated by the letters D (Depression), A (Anxiety) and S (Stress). For each scale (D, A & S) sum the scores for identified items. Because the DASS 21 is a short form version of the DASS (the Long Form has 42 items), the final score of each item groups (Depression, Anxiety and Stress) needs to be multiplied by two (x2).

Procedure

The data of the present study will be collected by the investigator through personal contact with the participants at their colleges. Before administering the questionnaire, the purpose of study will be explained and participants will be assured that their responses will be kept confidential and is used for research purpose only. The investigator established the rapport; the participants will be requested to provide the true responses on the test booklet. The participants will be required to follow the instructions given in the questionnaire which will help the participants to understand how he/she has to give responses in an appropriate manner. After completing the booklet were collected from the respondents for scoring.

The Relationship Between Overweight and Psychological Problems Among University Students in Saudi Arabia

Analysis of data

Data of present study will be analyzed using suitable statistical techniques. This includes descriptive (mean, median, mode, standard deviation, percentage etc.) and correlation and multivariate statistical method such as correlation and other methods relevant to the data of the present study.

RESULT AND INTERPRETATION

The result of the study will be interpreted in the light of the findings of the study. The obtained result would then be discussed in the light of the hypothesis framed for the study and available literature. Conclusion will be drawn and suggestions as well recommendations will be proposed accordingly.

Table 1: Coefficient of Correlation of Scores on body Mass Index and different dimensions of Psychological well-being

Variables	Mean	SD	BMI	Anxiety	Depression	Stress	Overall
Body Mass Index	.79	1.04	1.00				
Depression	7.01	4.14	.61**	1.00			
Anxiety	6.50	4.28	.78**	.79**	1.00		
Stress	8.90	3.97	.63**	.84**	.87**	1.00	
Overall	17.50	12.62	.66**	.89**	.95**	.98**	1.00

** Correlation is significant at the 0.01 level

Correlational Analysis

Pearson's coefficient of correlation was used to examine the relationship between scores on body Mass Index and different dimensions of Psychological well-being. Results presented in Table 1 indicate significant positive relationship between body mass index and depression aspect of psychological wellbeing ($r = .61, p < .01$). This indicates that participants who are having high body mass index reported more depression. Body mass index was found significantly and positively correlated with anxiety aspect of psychological wellbeing ($r = .78, p < .01$). This means that participants who are having high body mass index reported more anxiety. Also, body mass index was significantly and positively correlated with stress aspect of psychological wellbeing ($r = .63, p < .01$). This indicates that participants who are having high body mass index reported poor psychological wellbeing.

Body mass index was found significantly and positively correlated with overall score of psychological wellbeing ($r = .78, p < .01$). This means that participants who are having high body mass index reported more anxiety.

Implications of the study

Despite the fact that many investigations confirm the importance of psychological factors in the development of obesity; these findings are frequently underestimated in prevention and treatment of obesity. Considering the high prevalence and increasing incidence of obesity in Saudi Arabia, which affect all age groups, it is necessary to implement adequate health promotion programs. Colleges provide a good setting to develop such interventions, which should comprise evaluation of students' health status, including mental health and nutritional aspects. It is also important to take into account that unhealthy behavior could affect self-esteem and educational achievement.

CONCLUSION

This research will also provide an overview of current evidence on the relationship between overweight and mental health for university students in Saudi Arabia. It will draw particular attention to the risk factors associated with overweight and mental health. This will also addresses issues around inequalities, the implications of psychological distress caused by weight related stigma and discrimination. It also highlights specific issues relating to University students overweight and summarizes current knowledge on the effectiveness of interventions, best practice and scope for services for those at risk.

REFERENCES

- Ajilore O, Haroon E, Kumaran S, Darwin C, Binesh N, Mintz J, Miller J, Thomas M. A, Kumar A. (2007). Measurement of brain metabolites in patients with type 2 diabetes and major depression using proton magnetic resonance spectroscopy. *Neuropsychopharmacology* 2007, 32:1224-1231.
- Al-Hazzaa H. M, Musaiger A. O, (2011). ATLS Research Group. Arab teens lifestyle study (ATLS): objectives, design, methodology and implications. *Diabetes Metab Syndr Obes*; 4: 417-426.
- Allison D. B, Mentore J. L, Heo M, Chandler L. P, Cappelleri J. C, Infante M. C, Weiden PJ. (1999): Antipsychotic-induced weight gain: a comprehensive research synthesis. *Am J Psychiatry*, 156:1686-1696.
- Al-Othaimen A.I, Al-Nozah M, Osman A.K. (2007). Obesity: an emerging problem in Saudi Arabia. Analysis of data from the national nutrition survey. *East Mediter Health J*; 13(2): 411-418.
- Al-Rethaiaan A. S, Fahmy A. A, Al-Shwaiyat N. M. (2010). Obesity and eating habits among college students in Saudi Arabia: a cross sectional study. *Nutr J*; 9: 39.
- de Wit L, Luppino F, van Straten A, Penninx B, Zitman F, Cuijpers P. (2010). Depression and obesity: a meta-analysis of community-based studies. *Psychiatry Res*, 178:230-235.
- disorders in the population: a systematic review and meta-analysis. *International Journal of Obesity*; 34:407-19.
- Fernandes J, Arts J, Diamnd E, Hirshberg S, Lofgren I. E. (2013). Dietary factors are associated with coronary heart disease risk factors in severe obesity. *Nutr Res*; 33(8): 647-652.
- Gadalla T, Piran N (2008). Psychiatric comorbidity in women with disordered eating behavior: a national study. *Women Health*, 48:467-484.
- Garipey G, Nitka D, Schmitz N. (2010). The association between obesity and anxiety
- Heva A, Laitinen J, Miettunen J, et al. (2006). Obesity and depression: results from the longitudinal Northern Finland 1966 Birth Cohort Study. *Int J Obes.*;30(3):520-527.
- Iannotti R. J, Wang J. (2013). Trends in physical activity, sedentary behavior, diet, and BMI among US adolescents, 2001-2009. *Pediatrics* 2013 Oct; 132(4): 606-614.
- James P. T, Leach R, Kalamara E and Shayeghi M, "The worldwide obesity epidemic", *Obes Res.*, Vol 9 (Suppl 4), 2001, pp. 228S–233S.
- Khalaf A, Ekblom O, Kowalski J, Berggren V, Westergren A, Al- HazzaaH. (2013). Female university students' physical activity levels and associated factorsda cross-sectional study in South western Saudi Arabia. *Int J Environ Res Public Health*; 10: 3502e3517.
- Kivimäki M, Lawlor D. A, Singh-Manoux A, Batty G. D, Ferrie J. E, Shipley M.J, Nabi H, Sabi S, Marmot MG, Jokela M. (2009). Common mental disorder and obesity:

The Relationship Between Overweight and Psychological Problems Among University Students in Saudi Arabia

- insight from four repeat measures over 19 years: prospective Whitehall II cohort study. *BMJ*, 339: b3765.
- Luppino F. S, de Wit L. M, Bouvy P. F, Stijnen T, Cuijpers P, Penninx B, Zitman FG: (2010). Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Arch Gen Psychiatry*, 67: 220-229.
- Luppino F. S, de Wit LM, Bouvy P. F, Stijnen T, Cuijpers P, Penninx B, et al. (2010). Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Archives of General Psychiatry* 2010; 67(3):220-9.
- Mahfouz A.A, Abdelmoneim I, Khan M.Y, Daffalla A.A, Diab M.M, Al-Gelban K. S, et al. (2007). Obesity and related behaviors among adolescent school boys in Abha City, Southwestern Saudi Arabia. *J Trop Pediatr*; 54(2): 120e124.
- Needham B. L., Epel E. S, Adler N. E, Kiefe C. (2010). Trajectories of Change in Obesity and Symptoms of Depression: The CARDIA Study. *Am J Public Health* 2010, 100:1040-1046.
- Norah M. (2010). Obesity among Saudi female university students: dietary habits and health behaviors. *J Egypt Public Health Assoc.* 85(1): 46-59.
- Richardson L. P, Davis R, Poulton R, McCauley E, Moffitt T. E, Caspi A, Connell F: A (2003). Longitudinal evaluation of adolescent depression and adult obesity. *Arch Pediatr Adolesc Med*, 157:739-745.
- Stunkard AJ, Faith MS, Allison KC. (2003). Depression and obesity. *Biol Psychiatry*. 54, 330-337
- Swallen K. C, Reither E. N, Haas S. A, Meier A. M. (2005). Overweight, obesity, and health-related quality of life among adolescents: the National Longitudinal Study of Adolescent Health. *Pediatrics*. 115:340-347.
- Walker B. R. (2001). Activation of the hypothalamic-pituitary-adrenal axis in obesity: Cause or consequence? *Growth Hormone & IGF Research*, 11(Suppl A): S91-S95.
- WHO. Global recommendations on physical activity for health (2010). Available online: <http://www.who.int/dietphysicalactivity/global-PA-recs-2010.pdf>; [accessed on 15.06.14].
- World Federation for Mental Health, (2010). Mental Health and chronic physical illnesses: The need for continued and integrated care. <http://wfmh.com/wp-content/uploads/2014/02/WMHDAY2010.pdf>.
- Yaguchi-Tanaka Y, Kawagoshi Y, Sasaki S, Fukao A. (2013). Crosssectional study of possible association between rapid eating and high body fat rates among female Japanese college students. *J Nutr Sci Vitaminol*; 59(3): 243-249.

Acknowledgment

The author(s) appreciates all those who participated in the study and helped to facilitate the research process.

Conflict of Interest

The author(s) declared no conflict of interest.

How to cite this article: Pathath, A.W. & Begum, N. (2024). The Relationship Between Overweight and Psychological Problems Among University Students in Saudi Arabia. *International Journal of Indian Psychology*, 12(2), 029-035. DIP:18.01.004.20241202, DOI:10.25215/1202.004