

## Antisocial Personality Disorder, It's Risk Factors and Treatment

Ms. Pooja Gahlot<sup>1\*</sup>, Mr. Aniket Biraje<sup>2</sup>

### ABSTRACT

Personality disorders are mental health conditions characterized by a lifelong pattern of self-perception and problematic responses to others. Individuals who suffer from personality disorders frequently struggle to accept and understand their feelings. This study aims to provide a thorough overview of Antisocial Personality Disorder (ASPD), a subtype of Personality Disorder. It is a review paper. It explores prevalence, critical traits, diagnostic classifications, and related risk factors. The review covers a range of therapeutic modalities, such as Mentalization-Based Therapy, Cognitive Behaviour Therapy, Schema Therapy, and the Risk-Need-Responsivity model. The study highlights the difficulties in treating personality disorders as well as the critical role that each patient's motivation plays in achieving positive therapeutic outcomes.

**Keywords:** *Antisocial Personality Disorder, Risk Factors, Treatment*

Personality disorders are patterns from domains such as cognitive, emotional, control of impulses, and interpersonal interactions; they are non-functional, disrupted, and persistent over time, making it difficult to function in day-to-day living. Cluster A, Cluster B, Cluster C, and Other Personality Disorders are the four groups into which personality disorders are divided in the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5). Clusters A, B, and C are generally characterized by (1) odd and eccentric qualities; (2) emotional, dramatic, and erroneous traits; and (3) anxious and fearful actions, in that order. However, the World Health Organisation (WHO) has replaced its prior categorical classification system with a dimensional one that focuses on the severity of impairments to social and self-functioning abilities in the International Classification of Diseases (ICD-11). The ICD-11 categorizes personality disorders into four categories: (1) Mild Personality Disorder (2) Moderate Personality Disorder; (3) Severe Personality Disorder; and (4) Personality Disorder with unspecified severity. In addition to the severity of the personality disorders, traits domains such as anankastia, borderline pattern, detachment, disinhibition, dissociality, and negative affectivity are utilised to elucidate specific aspects. The DSM-5 Antisocial Personality Disorder and the ICD-11 Personality Disorders with Dissocial Traits shares similarities.

<sup>1</sup>Psychologist at Abhasa Rehabilitation Centre, Mumbai Maharashtra.

<sup>2</sup>Assistant Professor at Department of Psychology, Samarth Educational Institute, Dervan, Chiplun, Maharashtra.

\*Corresponding Author

Received: January 24, 2024; Revision Received: April 12, 2024; Accepted: April 16, 2024

## Antisocial Personality Disorder, It's Risk Factors and Treatment

Not until the DSM-III was published in 1980 did the term antisocial personality disorder even make an appearance in the psychiatric field. People frequently use the terms sociopathy and psychopathy interchangeably in conversation since they both seem to characterize someone who lacks regret. While there are certain similarities between sociopathy and psychopathy, experts continue to dispute some significant distinctions that lead to two distinct disorders and diagnoses rather than just one.

The hallmarks of antisocial personality disorder, also known as dissocial personality disorder, include an inconsiderate disregard for social obligations and other people's feelings. There is a glaring discrepancy between the behaviour and accepted societal norms. Behaviour cannot easily be changed by unfavorable experiences, such as punishment. There is a propensity to place blame on others or provide gratifying justifications for the activity that puts the patient at odds with society. There is also a low threshold for aggression, including violence, and minimal tolerance for frustration.

The only disorder among the Cluster B personality disorders that cannot be diagnosed in children is antisocial personality disorder. Early onset of ASPD—typically by the age of eight—is accompanied by a diagnosis of conduct disorder in childhood. If antisocial behaviour persists, the diagnosis is changed to ASPD by age eighteen. Co-occurring mental health, addiction, and medical comorbidities are linked to ASPD.

In a population sample of 350 emerging people, a study was done to investigate potential interactions between impulsivity, empathic issues, family functioning, and antisocial personality disorders. The findings indicated that motor impulsivity, sympathetic concern, and parental behaviour control were predicted factors for antisocial personality problems. High rates of comorbidity and mortality for persons with ASPD, along with detrimental effects on their family and society, characterise this complex disorder. People with ASPD typically perform poorly on socialization tests. This disease draws attention to the stark differences in behaviour between those who exhibit self-importance and an inability to love and adhere to social rules. These people are callous, inconsiderate, and reckless. Compared to people without the disorders, people with ASPD are reported to communicate poorly, confide less in their partners, and experience more emotional abuse in their relationships. As a result, having ASPD has a severe detrimental impact on a person's life as well as the lives of their friends, family, and the larger society.

According to estimates, ASPD affects 0.6% to 3.6% of adults. Men are three times more likely than women to have it, and among male inmates, it may rise to 60%. Medical and psychological comorbidities are more common in older persons with ASPD.

In addition to each physical disease, older persons with ASPD are more likely to receive a diagnosis of a substance use disorder, significant depression, mania, and generalized anxiety disorder.

Childhood emotional and physical abuse are often comorbid with ASPD. ASPD and physical neglect are significantly correlated. Children who experience abuse or neglect and poor parental connection are more likely to develop ASPD. Additionally, there is a strong link between ASPD sufferers and overprotective parents.

Many individuals with ASPD diagnoses experience difficulties interacting with others and wind up in the criminal justice system. The worldview of antisocial people is personal rather

## **Antisocial Personality Disorder, It's Risk Factors and Treatment**

than relational. They are unable to put another person's viewpoint above their own. They are unable to assume the position of somebody else as a result.

### ***Risk Factors***

There is no known cause of ASPD. Both environmental and genetic factors—such as child abuse—have a role in the development of this illness. Individuals who have an alcoholic or antisocial parent are more vulnerable. Increased adverse childhood experiences are thought to be linked to an ASPD diagnosis, with physical abuse demonstrating correlations with ASPD symptoms and sexual abuse linked to an ASPD diagnosis over the course of a lifetime.

Based on a 2019 study, 38–60% of ASPD diagnoses are related to hereditary factors. Growing up in a society with limited resources, being among peers who conduct in an antisocial manner, and seeing acts of social violence are all linked to social risk factors for the emergence of ASPD.

Risk factors for pregnancy and childbirth include being exposed to heavy metals, stress and anxiety throughout the pregnancy, smoking or using drugs during pregnancy, and problems during delivery.

A greater risk of developing ASPD also exists for children who are exposed to criminal activity and brutality as normal ways of life. The absence of brain regions linked to reasoning, empathy, and problem-solving in children who did not receive appropriate brain development may also put them at danger.

An increasing number of studies utilising brain imaging have demonstrated both structural and functional deficits in violent and antisocial people. The prefrontal cortex (especially the orbitofrontal and dorsolateral prefrontal cortex), the superior temporal gyrus, the amygdala hippocampus complex, and the anterior cingulate cortex are important areas that are frequently reported to be impaired in antisocial populations. In a study including twenty ASPD patients, researchers discovered that the hippocampal and amygdala volumes on both sides of antisocial personality disorder patients were statistically considerably lower than those of healthy control participants.

One neurotransmitter that controls aggression, mood, and social conduct is serotonin (5-HT). Research has demonstrated a connection between ASPD and serotonergic genes. Dopamine release is higher in ASPD patients than in healthy people. According to a study, those with high levels of ASPD characteristics emitted up to four times as much dopamine. Evidence suggests that extremely stressful daily settings are a major factor in the development of antisocial conduct in children from low-income families. Understanding more about how stressful daily circumstances impact low-income children's day-to-day adjustment and risk for antisocial conduct could be a potential approach to improving the lives of these children. Therefore, it appears that early exposure to abuse, socioeconomic position, and a lawless community are important environmental factors in the development of ASPD symptoms. The results of a study indicated that a range of executive functioning deficiencies, or deficiencies in cognitive control, are linked to symptoms of ASPD.

### ***Treatment***

Like other personality disorders, antisocial personality disorder is extremely challenging to cure. Convincing an individual that they require treatment is the first obstacle. The Food and

## Antisocial Personality Disorder, It's Risk Factors and Treatment

Drug Administration (FDA) has not approved any medications expressly to treat ASPD. Medical professionals may recommend medication for conditions including anxiety, depression, or aggressive symptoms that can occasionally coexist with ASPD. Mood stabilisers, which assist in managing abrupt changes in mood or behaviour, and antipsychotics, which control aggressive or violent behaviour, are advised. According to studies, some ASPD sufferers found success with Cognitive Behaviour Therapy, which tries to alter dysfunctional thought and behaviour patterns. "Like most mental health disorders, the desire for change must come from the person"—that is, psychotherapy for ASPD can only be effective if the person receiving it is truly motivated to change. The overall consensus in the research is that CBT is a useful therapeutic approach for patients with personality disorders, as it reduces symptoms and improves functional outcomes. The ASPD is one of the personality disorders that are conceptualised by the cognitive theory of personality disorders based on their fundamental beliefs or schemas. Antisocial patients have an extremely negative perception of other people; they believe that other people have been exploited and should therefore also be exploited in retribution. People who suffer from personality disorders tend to have strong negative core beliefs and extremely weak positive core beliefs. Usually, it is the unfavourable underlying belief that rules their existence. CBT methods assist a person in altering their fundamental beliefs. The research' findings indicate that while non-significant behavioural and attitudinal improvements are the result of typical CBT treatments, a new cognitive behavioural model known as the risk-need responsibility (RNR) has emerged as a promising treatment for ASPD.

The three core RNR principles are as follows: Risk is the decision of whom to target based on a person's propensity for future offences. This is significant because the type of intervention should correspond to the chance of reoffending; low-risk patients should receive treatment, while high-risk cases should receive rehabilitation obtaining minimum assistance. Need is about what ought to be done; targeted treatments should centre on identified criminogenic needs rather than other needs unrelated to criminal conduct. Delivering the work in a responsible manner involves both general and specialised responsivity. Specific responsivity states that interventions should be customised to the individual's abilities among other things, while general responsivity encourages the use of cognitive social learning techniques to impact behaviour. Responsivity includes the ability to supervise.

For people with ASPD, routine maintenance combined with contingency management is also claimed to be beneficial. In a sort of behaviour therapy known as "contingency management," people receive praise or rewards when they demonstrate a change in their behaviour. People with ASPD may have better social functioning with contingency management.

Based on the principles of schema theory, Schema Therapy has become a popular therapy option for personality disorders. Although it incorporates aspects of both object relations theory and attachment theory, cognitive behavioural theory serves as the primary foundation for schema therapy. People with ASPD frequently have inflexible, maladaptive schemas that are linked to emotional repression, mistrust/abuse, anger, and unfulfilled needs from their early years. In order to establish a therapeutic rapport with the client, Schema Therapy employs two strategies: limited reparenting and empathic confrontation. Limited reparenting involves showing empathy on the side of the psychologist; in contrast, empathic confrontation entails striking a balance between demonstrating empathy and helping the client see how their maladaptive schemas have unfavourable effects. In order to improve their coping

## Antisocial Personality Disorder, It's Risk Factors and Treatment

mechanisms and adopt more acceptable and healthy behaviours, participants in this therapy must discover the underlying causes of their behaviour.

Another talking therapy that's gaining momentum in the management of ASPD is mentalization-based therapy (MBT). With MBT, the therapist will prompt the patient to think about their thoughts and how their mental health impacts their actions. The capacity to think about thinking is known as mentalization. This entails analysing your own ideas and convictions to determine whether they are practical, grounded in reality, and realistic. Four overlapping aspects can be used to describe mentalization: self/other oriented, automatic/controlled, implicit/explicit, internally/externally based, and cognitive/affective process. Family therapy can potentially mitigate the negative effects of ASPD on an individual and the family system by promoting healthy boundaries, mutual understanding, and communication among individuals with ASPD and their families.

### CONCLUSION

Personality disorders are persistent patterns in cognitive, emotional, impulse control, and interpersonal interactions. Antisocial personality disorder (ASPD) is the only disorder in Cluster B that cannot be diagnosed in children and is characterized by disregard for social obligations, discrepancy between behavior and societal norms, propensity to blame others, low aggression threshold, and minimal tolerance for frustration. It affects 0.6%-3.6% of adults, with men being three times more likely to have it. ASPD is linked to environmental and genetic factors, with early exposure to abuse, socioeconomic position, and a lawless community contributing to 38-60% of diagnoses. Brain imaging studies show structural and functional deficits in areas like the prefrontal cortex, superior temporal gyrus, amygdala hippocampus complex, and anterior cingulate cortex. Cognitive behavior therapy (CBT) and risk-need responsibility (RNR), MBT, Schema Therapy treatments are effective in treating ASPD, as they reduce symptoms and improve functional outcomes.

### REFERENCES

- Antisocial personality disorder – Diagnosis and treatment – Mayo Clinic. (2023, February 24). <https://www.mayoclinic.org/diseases-conditions/antisocial-personality-disorder/diagnosis-treatment/drc-20353934#:~:text=There%20are%20no%20medicines%20specifically,or%20for%20symptoms%20of%20aggression>
- Antisocial Personality Disorder Symptoms, Causes & Treatment. (n.d.). Sheppard Pratt. <https://www.sheppardpratt.org/knowledge-center/conditions-diagnoses/antisocial-personality-disorder/Antisocial>
- Antisocial Personality Disorder Treatment. (2023, December 29). Choosing Therapy. <https://www.choosingtherapy.com/antisocial-personality-disorder-treatment/>
- Antisocial Personality Disorder. (2023, September 9). Annabelle Psychology | Singapore's Leading Psychologists. <https://www.annabellepsychology.com/antisocial-personality-disorder#:~:text=Schema%20therapy,-While%20it%20is&text=Individuals%20with%20ASPD%20tend%20to,unmet%20needs%20in%20early%20childhood>
- Antisocial personality disorder. (2024, January 6). Wikipedia. [https://en.m.wikipedia.org/wiki/Antisocial\\_personality\\_disorder#:~:text=ASPD%20is%20highly%20comorbid%20with,early%20in%20life%20is%20importa](https://en.m.wikipedia.org/wiki/Antisocial_personality_disorder#:~:text=ASPD%20is%20highly%20comorbid%20with,early%20in%20life%20is%20importa)
- Antisocial Personality Disorder. (n.d.). SAMHSA. <https://www.samhsa.gov/mental-health/a/antisocial-personality-disorder#:~:text=Causes,parent%20are%20at%20increased%20risk>

## Antisocial Personality Disorder, It's Risk Factors and Treatment

- Antisocial Personality Disorder: Often Overlooked and Untreated. (n.d.). <https://www.psychiatry.org/news-room/apa-blogs/antisocial-personality-disorder-often-overlooked#:~:text=It%20is%20estimated%20to%20affect,common%20among%20men%20than%20women.&text=Individuals%20with%20antisocial%20personality%20disorder,disorder%20or%20substance%20use%20disordersAntisocial>
- Antisocial Personality Disorder: Symptoms, Causes, &#038; Treatments. (2023, December 29). Choosing Therapy. <https://www.choosingtherapy.com/antisocial-personality-disorder/Antisocial>
- Are people-pleasing and sociopathy opposite ends of the same spectrum ... Available at: <https://www.psychologytoday.com/us/blog/healthy-brain-happy-life/202110/are-people-pleasing-and-sociopathy-opposite-ends-the-same>.
- Bateman, A., Bolton, R., & Fonagy, P. (2013, January 1). Antisocial Personality Disorder: A Mentalizing Framework. *Focus*. <https://doi.org/10.1176/appi.focus.11.2.178>
- Bell, M.C. (2009) Antisocial Personality Disorder and Romantic Relationship Functioning in an Epidemiologically-Based Sample.
- Black, D. W. (2015, July 1). The Natural History of Antisocial Personality Disorder. *The Canadian Journal of Psychiatry*. <https://doi.org/10.1177/070674371506000703Black>
- Black, D.W. (2015) The natural history of antisocial personality disorder, *Canadian journal of psychiatry*. *Revue canadienne de psychiatrie*. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4500180/>.
- DeLisi, M., Drury, A. J., & Elbert, M. J. (2019, July 1). The etiology of antisocial personality disorder: The differential roles of adverse childhood experiences and childhood psychopathology. *Comprehensive Psychiatry*. <https://doi.org/10.1016/j.comppsy.2019.04.001>
- Environmental Effect on Antisocial Behavior Disorder | Psychology Paper Example. (2023, September 22). PsychologyWriting. <https://psychologywriting.com/environmental-effect-on-antisocial-behavior-disorder/>
- Farias, M.S.J.A. de et al. (2013) The Antisocial Personality Disorder and its characteristics, changes and advances: a challenge, *International Journal of Collaborative Research on Internal Medicine and Public Health*.
- Fisher, K. A. (2023, May 8). Antisocial Personality Disorder. *StatPearls – NCBI Bookshelf*. <https://www.ncbi.nlm.nih.gov/books/NBK546673/Fisher>
- Gibbon, S., Duggan, C., Stoffers, J. M., Huband, N., Völlm, B., Ferriter, M., & Lieb, K. (2010, June 16). Psychological interventions for antisocial personality disorder. *The Cochrane Library*. <https://doi.org/10.1002/14651858.cd007668.pub2>
- Haelle, T. (2022, October 12). How to Treat Antisocial Personality Disorder. *Everyday Health.com*. <https://www.everydayhealth.com/antisocial-personality-disorder/treatment/>
- Haelle, T. (2022, October 23). Complications and Life Consequences of Antisocial Personality Disorder. Retrieved from <https://www.everydayhealth.com/antisocial-personality-disorder/life-consequences/Haelle>
- Houser, M. C. (2015, April 1). A History of Antisocial Personality Disorder in The Diagnostic and Statistical Manual of Mental Illness and Treatment from A Rehabilitation Perspective. *CORE*. <https://core.ac.uk/works/33699272>
- Houser, M.C. (2015) A History of Antisocial Personality Disorder in the Diagnostic and Statistical Manual of Mental Illness and Treatment from A Rehabilitation Perspective, *OpenSIUC*.
- ICD-10 Version:2016. (n.d.). <https://icd.who.int/browse10/2016/en#F60.2ICD-10>
- Impulsive-Antisocial Personality Traits Linked to a Hypersensitive. (2015, August 12). National Institutes of Health (NIH). <https://www.nih.gov/news-events/news->

## Antisocial Personality Disorder, It's Risk Factors and Treatment

- releases/impulsive-antisocial-personality-traits-linked-hypersensitive-brain-reward-system#:~:text=%22The%20amount%20of%20dopamine%20released,lead%20author%20of%20the%20studymple%2C%20one%20research%20group,a%20more%20robust%20dopamine%20system
- Kaya, H., Yildirim, H., & Atmaca, M. (2020, May 1). Reduced hippocampus and amygdala volumes in antisocial personality disorder. *Journal of Clinical Neuroscience*. <https://doi.org/10.1016/j.jocn.2020.01.048>
- Lautieri, A. (2019, March 25). Schema Therapy for Personality Disorders. *MentalHelp.net*. <https://www.mentalhelp.net/personality-disorders/schema-therapy/>
- Lovering, C. (2021, December 23). Causes of Antisocial Personality Disorder. *Psych Central*. <https://psychcentral.com/disorders/antisocial-personality-disorder/antisocial-personality-disorder-causes#possible-causes>
- Marzilli, E., Cerniglia, L., & Cimino, S. (2021, May 23). Antisocial Personality Problems in Emerging Adulthood: The Role of Family Functioning, Impulsivity, and Empathy. <https://doi.org/10.3390/brainsci11060687Marzilli>
- Mateo, A. (2022, June 30). What To Know About Antisocial Personality Disorder Treatment. *Health*. <https://www.health.com/condition/antisocial-personality-disorder/antisocial-personality-disorder-treatment>
- Matusiewicz, A.K. et al. (2010) The effectiveness of cognitive behavioral therapy for personality disorders, *The Psychiatric clinics of North America*. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3138327/>.
- Michael A. Russell, M.A., Candice L. Odgers, Ph.D , Antisocial Behavior among Children in Poverty: Understanding Environmental Effects in Daily Life, Environmental Effects on Antisocial Behavior.
- Moran, P. (1999, May 27). The epidemiology of antisocial personality disorder. *Social Psychiatry and Psychiatric Epidemiology*. <https://doi.org/10.1007/s001270050138Moran>
- Navigating Antisocial Personality Disorder: Family Therapy | *BetterHelp*. (2024, January 6). <https://www.betterhelp.com/advice/personality-disorders/navigating-antisocial-personality-disorder-family-therapy/>
- Petry, N.M. (2011) Contingency management: What it is and why psychiatrists should want to use it, *The psychiatrist*. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3083448/#:~:text=Contingency%20management%20refers%20to%20a,evidence%20of%20positive%20behavioural%20change>.
- Prevalence and correlates of antisocial personality disorder in older ... Available at: <https://www.tandfonline.com/doi/full/10.1080/13607863.2020.1839867>.
- Professional, C. C. M. (n.d.). Antisocial Personality Disorder (ASPD). *Cleveland Clinic*. <https://my.clevelandclinic.org/health/diseases/9657-antisocial-personality-disorder>
- Sargin, A. E., Özdel, K., & Türkçapar, M. H. (2017, December 20). Cognitive-Behavioral Theory and Treatment of Antisocial Personality Disorder. *InTech eBooks*. <https://doi.org/10.5772/intechopen.68986>
- Sargin, A. E., Özdel, K., & Türkçapar, M. H. (2017, December 20). Cognitive-Behavioral Theory and Treatment of Antisocial Personality Disorder. *InTech eBooks*. <https://doi.org/10.5772/intechopen.68986>
- Sargin, A.E., Ozdel, K. and Turkcapar, M.H. (2017) Cognitive- Behavioral Theory and Treatment of Antisocial Personality Disorder, *INTECH*. Available at: <http://dx.doi.org/10.5772/intechopen.68986>.
- Simona, T. (2019, December 11). Antisocial Personality Disorder. Retrieved from [https://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=3502191Simona](https://papers.ssrn.com/sol3/papers.cfm?abstract_id=3502191Simona)

## Antisocial Personality Disorder, It's Risk Factors and Treatment

- The Risk-Need-Responsivity Model. (n.d.). <https://www.justiceinspectorates.gov.uk/hmiprobation/research/the-evidence-base-probation/models-and-principles/the-rnr-mode>
- Website, N. (2023, February 15). Treatment – Borderline personality disorder. Nhs.uk. [https://www.nhs.uk/mental-health/conditions/borderline-personality-disorder/treatment/#:~:text=Mentalisation%2Dbased%20therapy%20\(MBT\)&text=MBT%20is%20based%20on%20the,realistic%20and%20based%20on%20reality](https://www.nhs.uk/mental-health/conditions/borderline-personality-disorder/treatment/#:~:text=Mentalisation%2Dbased%20therapy%20(MBT)&text=MBT%20is%20based%20on%20the,realistic%20and%20based%20on%20reality)
- Wong, R. S. Y. (2023, August 16). Psychopathology of antisocial personality disorder: from the structural, functional and biochemical perspectives. *The Egyptian Journal of Neurology, Psychiatry and Neurosurgery*. <https://doi.org/10.1186/s41983-023-00717-4>
- Wong, R. S. Y. (2023, August 16). Psychopathology of antisocial personality disorder: from the structural, functional and biochemical perspectives. *The Egyptian Journal of Neurology, Psychiatry and Neurosurgery*. <https://doi.org/10.1186/s41983-023-00717-4>
- Yakeley, J., & Williams, A. (2014, March 1). Antisocial personality disorder: new directions. <https://doi.org/10.1192/apt.bp.113.011205> Yakeley
- Yang, Y., Glenn, A. L., & Raine, A. (2008, January 1). Brain abnormalities in antisocial individuals: implications for the law. *Behavioral Sciences & the Law*. <https://doi.org/10.1002/bsl.788>
- Zeier JD, Baskin-Sommers AR, Hiatt Racer KD, Newman JP. Cognitive control deficits associated with antisocial personality disorder and psychopathy. *Personal Disorder*. 2012 Jul;3(3):283-93. Doi: 10.1037/a0023137. Epub 2011 Jul 4. PMID: 22452754; PMCID: PMC3387332.

### **Acknowledgment**

The author(s) appreciates all those who participated in the study and helped to facilitate the research process.

### **Conflict of Interest**

The author(s) declared no conflict of interest.

**How to cite this article:** Gahlot, P. & Biraje, A. (2024). Antisocial Personality Disorder, It's Risk Factors and Treatment. *International Journal of Indian Psychology*, 12(2), 159-166. DIP:18.01.017.20241202, DOI:10.25215/1202.017